



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/53, 15/82, 9/02, A01H 5/00	A2	(11) International Publication Number: WO 96/13591 (43) International Publication Date: 9 May 1996 (09.05.96)
(21) International Application Number: PCT/US95/13784 (22) International Filing Date: 25 October 1995 (25.10.95) (30) Priority Data: 08/329,560 26 October 1994 (26.10.94) US (71) Applicant: MICHIGAN STATE UNIVERSITY [US/US]; 238 Administration Building, East Lansing, MI 48824 (US). (72) Inventors: CAHOON, Edgar, B.; Building 463, Brookhaven National Laboratories, Upton, NY 11973 (US). OHLRO- GEE, John, B.; 2123 Lagoon Drive, Okemos, MI 48864 (US). (74) Agents: YATES, Michael, E. et al.; 700 Capital Square, 400 Locust Street, Des Moines, IA 50309 (US).	(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, LS, MW, SD, SZ, UG). Published Without international search report and to be republished upon receipt of that report.	
(54) Title: METHODS AND COMPOSITIONS RELATING TO PLANT Δ^6 PALMITOYL-ACYL CARRIER PROTEIN DESATURASE		
(57) Abstract <p>A plant Δ^6 palmitoyl-acyl carrier protein desaturase, the gene encoding the desaturase, and transgenic plants and plant cells containing the heterologous DNA encoding the desaturase are described. The desaturase introduces a double bond at the sixth carbon atom from the carboxyl end of a 16 carbon saturated fatty acid, and is therefore useful in production of plant seeds having a modified fatty acid profile.</p> <pre> 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 1075 1076 1077 1078 1079 1080 1081 1082 1083 1084 1085 1086 1087 1088 1089 1090 1091 1092 1093 1094 1095 1096 1097 1098 1099 1100 1101 1102 1103 1104 1105 1106 1107 1108 1109 1110 1111 1112 1113 1114 1115 1116 1117 1118 1119 1120 1121 1122 1123 1124 1125 1126 1127 1128 1129 1130 1131 1132 1133 1134 1135 1136 1137 1138 1139 1140 1141 1142 1143 1144 1145 1146 1147 1148 1149 1150 1151 1152 1153 1154 1155 1156 1157 1158 1159 1160 1161 1162 1163 1164 1165 1166 1167 1168 1169 1170 1171 1172 1173 1174 1175 1176 1177 1178 1179 1180 1181 1182 1183 1184 1185 1186 1187 1188 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1200 1201 1202 1203 1204 1205 1206 1207 1208 1209 1210 1211 1212 1213 1214 1215 1216 1217 1218 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1230 1231 1232 1233 1234 1235 1236 1237 1238 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1260 1261 1262 1263 1264 1265 1266 1267 1268 1269 1270 1271 1272 1273 1274 1275 1276 1277 1278 1279 1280 1281 1282 1283 1284 1285 1286 1287 1288 1289 1290 1291 1292 1293 1294 1295 1296 1297 1298 1299 1300 1301 1302 1303 1304 1305 1306 1307 1308 1309 1310 1311 1312 1313 1314 1315 1316 1317 1318 1319 1320 1321 1322 1323 1324 1325 1326 1327 1328 1329 1330 1331 1332 1333 1334 1335 1336 1337 1338 1339 1340 1341 1342 1343 1344 1345 1346 1347 1348 1349 1350 1351 1352 1353 1354 1355 1356 1357 1358 1359 1360 1361 1362 1363 1364 1365 1366 1367 1368 1369 1370 1371 1372 1373 1374 1375 1376 1377 1378 1379 1380 1381 1382 1383 1384 1385 1386 1387 1388 1389 1390 1391 1392 1393 1394 1395 1396 1397 1398 1399 1400 1401 1402 1403 1404 1405 1406 1407 1408 1409 1410 1411 1412 1413 1414 1415 1416 1417 1418 1419 1420 1421 1422 1423 1424 1425 1426 1427 1428 1429 1430 1431 1432 1433 1434 1435 1436 1437 1438 1439 1440 1441 1442 1443 1444 1445 1446 1447 1448 1449 1450 1451 1452 1453 1454 1455 1456 1457 1458 1459 1460 1461 1462 1463 1464 1465 1466 1467 1468 1469 1470 1471 1472 1473 1474 1475 1476 1477 1478 1479 1480 1481 1482 1483 1484 1485 1486 1487 1488 1489 1490 1491 1492 1493 1494 1495 1496 1497 1498 1499 1500 1501 1502 1503 1504 1505 1506 1507 1508 1509 1510 1511 1512 1513 1514 1515 1516 1517 1518 1519 1520 1521 1522 1523 1524 1525 1526 1527 1528 1529 1530 1531 1532 1533 1534 1535 1536 1537 1538 1539 1540 1541 1542 1543 1544 1545 1546 1547 1548 1549 1550 1551 1552 1553 1554 1555 1556 1557 1558 1559 1560 1561 1562 1563 1564 1565 1566 1567 1568 1569 1570 1571 1572 1573 1574 1575 1576 1577 1578 1579 1580 1581 1582 1583 1584 1585 1586 1587 1588 1589 1590 1591 1592 1593 1594 1595 1596 1597 1598 1599 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616 1617 1618 1619 1620 1621 1622 1623 1624 1625 1626 1627 1628 1629 1630 1631 1632 1633 1634 1635 1636 1637 1638 1639 1640 1641 1642 1643 1644 1645 1646 1647 1648 1649 1650 1651 1652 1653 1654 1655 1656 1657 1658 1659 1660 1661 1662 1663 1664 1665 1666 1667 1668 1669 1670 1671 1672 1673 1674 1675 1676 1677 1678 1679 1680 1681 1682 1683 1684 1685 1686 1687 1688 1689 1690 1691 1692 1693 1694 1695 1696 1697 1698 1699 1700 1701 1702 1703 1704 1705 1706 1707 1708 1709 1710 1711 1712 1713 1714 1715 1716 1717 1718 1719 1720 1721 1722 1723 1724 1725 1726 1727 1728 1729 1730 1731 1732 1733 1734 1735 1736 1737 1738 1739 1740 1741 1742 1743 1744 1745 1746 1747 1748 1749 1750 1751 1752 1753 1754 1755 1756 1757 1758 1759 1760 1761 1762 1763 1764 1765 1766 1767 1768 1769 1770 1771 1772 1773 1774 1775 1776 1777 1778 1779 1780 1781 1782 1783 1784 1785 1786 1787 1788 1789 1790 1791 1792 1793 1794 1795 1796 1797 1798 1799 1800 1801 1802 1803 1804 1805 1806 1807 1808 1809 1810 1811 1812 1813 1814 1815 1816 1817 1818 1819 1820 1821 1822 1823 1824 1825 1826 1827 1828 1829 1830 1831 1832 1833 1834 1835 1836 1837 1838 1839 1840 1841 1842 1843 1844 1845 1846 1847 1848 1849 1850 1851 1852 1853 1854 1855 1856 1857 1858 1859 1860 1861 1862 1863 1864 1865 1866 1867 1868 1869 1870 1871 1872 1873 1874 1875 1876 1877 1878 1879 1880 1881 1882 1883 1884 1885 1886 1887 1888 1889 1890 1891 1892 1893 1894 1895 1896 1897 1898 1899 1900 1901 1902 1903 1904 1905 1906 1907 1908 1909 1910 1911 1912 1913 1914 1915 1916 1917 1918 1919 1920 1921 1922 1923 1924 1925 1926 1927 1928 1929 1930 1931 1932 1933 1934 1935 1936 1937 1938 1939 1940 1941 1942 1943 1944 1945 1946 1947 1948 1949 1950 1951 1952 1953 1954 1955 1956 1957 1958 1959 1960 1961 1962 1963 1964 1965 1966 1967 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980 1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2</pre>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

METHODS AND COMPOSITIONS RELATING TO PLANT Δ^6 PALMITOYL-ACYL CARRIER PROTEIN DESATURASE

Field of the Invention

The present invention relates to plant fatty acid desaturases. More particularly, the present invention relates to plant Δ^6 palmitoyl-acyl carrier protein desaturases, the genes encoding such desaturases, transgenic plants and plant cells containing DNA encoding such desaturases, and methods for altering the fatty acid profile of plant seeds through the use of such desaturases.

BACKGROUND

Fatty acid desaturases of plants have received considerable attention because of their contributions to the physiology and economic value of plants. The activity of fatty acid desaturases, for example, may be a component of the ability of certain species to adjust levels of membrane unsaturation in response to stresses such as chilling (1-3). In addition, the degree of fatty acid unsaturation resulting from desaturase activity is often a major determinant of the nutritional and industrial quality of plant seed oils (4).

Plants typically contain a variety of fatty acid desaturases. The most numerous of these are membrane-associated desaturases that use fatty acids bound to glycerolipids as substrates (5). In addition, the synthesis of oleic acid (18:1 Δ^9) in plants and certain other organisms such as *Euglena* (photoauxotrophic) is catalyzed by a desaturase that functions on fatty acids esterified to acyl carrier protein (ACP) (6-8). This enzyme, the Δ^9 stearoyl (18:0)-ACP desaturase (EC 1.14.99.6), displays soluble activity in contrast to all previously characterized desaturases (6). In the presence of radiolabeled 18:0-ACP and cofactors including NADPH, ferredoxin (Fd), and ferredoxin-NADPH reductase, the activity of the Δ^9 18:0-ACP desaturase is readily detectable in extracts of most plant tissues (9). Due in part to its soluble nature and relative ease of assay, the Δ^9 18:0-ACP desaturase has been purified from several plant sources (10-14), and a number of cDNAs encoding

this enzyme have been isolated (12-20). In addition to the Δ^9 18:0-ACP desaturase, a Δ^4 16:0-ACP desaturase has recently been identified in plants (21, 22). This enzyme is a component of the petroselinic acid (18:1 Δ^6) biosynthetic pathway in endosperm of coriander (*Coriandrum sativum* L.) and other Umbelliferae species. Translation of a cDNA for the Δ^4 16:0-ACP desaturase has revealed that this enzyme shares extensive amino acid identity with the Δ^9 18:0-ACP desaturase (21).

The existence of structurally related acyl-ACP desaturases with different substrate recognition and double bond-positioning properties offers the opportunity to compare the active site structures of members of this family of enzymes using techniques such as site-directed mutagenesis and x-ray crystallography. Information gained from this research could potentially lead to the design of desaturases capable of producing new industrially useful isomers of monounsaturated fatty acids. These studies would be aided by the isolation of cDNAs for other variant acyl-ACP desaturases in addition to those for the Δ^9 18:0- and Δ^4 16:0-ACP desaturases. A potential source of such a desaturase is seed of *Thunbergia alata* (Acanthaceae family). The oil of this tissue consists of more than 80% weight of the unusual fatty acid Δ^6 hexadecenoic acid (16:1 Δ^6) (23). We have used biochemical and molecular biological approaches to examine whether 16:1 Δ^6 is synthesized by the activity of a unique acyl-ACP desaturase that is related to the Δ^9 18:0- and Δ^4 16:0-ACP desaturases.

SUMMARY

The present invention therefore relates to an isolated Δ^6 palmitoyl acyl carrier protein desaturase and DNA encoding same, transgenic plants and plant cells containing heterologous DNA encoding said desaturase, mRNA derived from DNA encoding such a desaturase, and vectors containing DNA encoding the desaturase. The present invention also relates to a method for modifying the fatty acid content of a plant seed involving the introduction of a double bond at the sixth carbon atom from the carboxyl end of the saturated 16 carbon fatty acid, for example palmitic acid. Additionally, the present

invention relates to plant seeds having a modified fatty acid content derived through the method of the invention.

An objective of the invention is to provide a mechanism for the modification of fatty acids in order to enhance the industrial and nutritional quality of plant seed oils. As part of this invention, we have developed a method for introducing a double bond at a novel position in a saturated fatty acid. The invention entails the identification of the enzyme (a Δ^6 -palmitoyl-acyl carrier protein desaturase) involved in the biosynthesis of the unusual fatty acid Δ^6 -hexadecenoic acid. In addition, a complementary DNA (cDNA) was isolated for this desaturase. Expression of the cDNA in *Escherichia coli* resulted in the production of a catalytically active Δ^6 -palmitoyl-acyl carrier protein desaturase.

This invention offers the opportunity to synthesize a monounsaturated fatty acid with a double bond positioned at the sixth carbon atom from the carboxyl end of the fatty acid molecule. The invention also allows for unsaturation to be introduced into a saturated 16 carbon fatty acid (palmitic acid). These two features of the invention allow for the production of a monounsaturated fatty acid that is not normally found in conventional plant seed oils. The major monounsaturated fatty acid present in seed oils is oleic acid, an 18 carbon fatty acid with a double bond positioned at the ninth carbon atom from the carboxyl end of the molecule.

The ability of the invention to introduce a double bond into palmitic acid offers the potential for reducing the saturated fatty acid content of vegetable oils. Typically, the presence of the 16 carbon saturated fatty acid palmitic acid limits the nutritional quality of seed oils. Therefore, the invention may allow for the production of a seed oil with reduced palmitic acid content. As a result, such an oil would presumably be less harmful to human health.

In addition, the product of the invention, Δ^6 -hexadecenoic acid, may be useful as a chemical precursor of certain industrial feedstocks. Because the double bond of Δ^6 -hexadecenoic acid is located at the sixth carbon atom, this fatty acid can be oxidatively cleaved to form adipic acid (a six carbon

dicarboxylic acid) and decanoic acid (a ten carbon fatty acid). Adipic acid is a precursor of nylon 66 and is currently derived from petroleum by-products.

Furthermore, the use of Δ^6 -hexadecenoic acid has been proposed as an emollient in cosmetics (U.S. Patent No. 4,036,991).

BRIEF DESCRIPTION OF THE DRAWINGS

FIGURE 1 is an autoradiogram of intact products of [^{14}C]16:0-ACP desaturase assays conducted with a 100,000 x g supernatant of *T. alata* endosperm extracts (16:1), and the methyl ester of 16:0 derived from unreacted [^{14}C]16:0-ACP(16:0). The Std. lane contains methyl [^{14}C]16:0 (16:0), [^{14}C]18:1 Δ^9 (Δ^9), and [^{14}C]18:1 Δ^6 (Δ^6).

FIGURE 2 is an autoradiogram of oxidized products of [^{14}C]16:0-ACP desaturase assays. Shown are the permanganate-periodate oxidation products of the methyl ester derivatives of 16:1 formed by acyl-ACP desaturation activity in *T. alata* endosperm homogenates. [^{14}C]16:0-ACP was used as the substrate for this assay.

FIGURE 3 shows the time course of [^{14}C]16:0-ACP desaturase activity in a 100,000 x g supernatant of a *T. alata* endosperm homogenate. Assays were conducted with 118 pmol of [^{14}C]16:0-ACP and 23 μg of protein.

FIGURE 4 shows the effect of catalase, ferredoxin (Fd), nitrogen (N_2), potassium cyanide (KCN), or hydrogen peroxide (H_2O_2) on Δ^6 16:0-ACP desaturase activity in *T. alata* endosperm extracts. Assays were conducted for 10 min using 118 pmol of [^{14}C]16:0-ACP and 23 μg of protein from a 100,000 x g supernatant of endosperm homogenate. Assays with potassium cyanide and hydrogen peroxide contained 1mM of each compound, and catalase was omitted from assays containing hydrogen peroxide. (n.d.—not detected).

FIGURE 5 is the nucleotide sequence of the cDNA insert of pTAD4 (Δ^6) and a comparison of the deduced amino acid sequences of pTAD4 (Δ^6) and cDNAs for the coriander Δ^4 16:0-ACP desaturase (Δ^4) (see Ref. 21) and the castor Δ^9 18:0-ACP desaturase (Δ^9) (see Ref. 13). Identical amino acids are indicated by colons. Amino acids that are absent relative to the castor Δ^9 18:0-ACP desaturase are indicated by dashed lines. Alignment of the

nucleotide sequence of the cDNA insert of pTAD4 is maintained with a dotted line. The underlined alanine at amino acid 33 is the likely start of the native peptide encoded by pTAD4.

FIGURE 6 compares the Δ^6 16:0-ACP desaturase activity of extracts of *E. coli* BL21 pLysS containing only the vector pET3d or pET3d with insert derived from the *T. alata* cDNA of pTAD4 (with or without isopropyl-1-thio- β -D-galactopyranoside induction). The methyl ester of 16:1 formed by the *E. coli*-expressed desaturase was separated from methyl 16:0 of the unreacted substrate by argentation TLC as shown. Assays were conducted for 60 min using 230 μ g of *E. coli* protein and 118 pmol of $[1-^{14}\text{C}]16:0$ -ACP. Shown in the standard chromatogram are (std.) are methyl $[^{14}\text{C}]16:0$, -18:1 Δ^9 , and -18:1 Δ^6 .

FIGURE 7 shows substrate specificity and ferredoxin dependence of the *E. coli*-expressed *T. alata* Δ^6 16:0-ACP desaturase encoded by the cDNA insert of pTAD4. Assays were performed for 10 minutes using 118 pmol of either $[1-^{14}\text{C}]14:0$ -, 16:0-(\pm ferredoxin, Fd), or 18:0-ACP and 65 μ g of total *E. coli* protein.

FIGURE 8 shows a mass spectrum of derivatives of $[1-^{14}\text{C}]16:1$ formed by 16:0-ACP desaturase activity in extracts of *E. coli* expressing the mature peptide-coding region of pTAD4. The $[1-^{14}\text{C}]16:1$ desaturation product was converted to a methyl ester derivative and reacted with dimethyl sulfoxide prior to mass spectral analysis.

DETAILED DESCRIPTION

Experimental Procedures

Plant Material - Studies were conducted using developing endosperm dissected from fruits of *T. alata* Bojer ex Sims (black-eyed susan vine) (Northrup King, Minneapolis, MN). Fruits were collected from plants grown either outdoors in pots during summers in East Lansing, Michigan or under greenhouse conditions with natural illumination. In the latter case, flowers required hand pollination for adequate fruit set. Endosperm was frozen in liquid nitrogen following dissection and stored at -70°C until use in enzyme assays or RNA extraction.

Acyl-ACP Desaturation Assays - Approximately 200-300 mg of developing *T. alata* endosperm was homogenized in 3 ml of buffer consisting of 100 mM Tris-HCl, pH 7.5, 2.5 mM dithiothreitol, 1 mM isoascorbate, 10% (v/v) glycerol, and 1.5% (w/v) polyvinylpolypyrrolidone using an Elvehjem tissue grinder. Debris and polyvinylpolypyrrolidone were subsequently removed by centrifugation at 14,000 x g for 5 min. The supernatant was then passed through two layers of miracloth (Calbiochem) and spun for an additional 10 min at 30,000 x g. The soluble phase was removed while attempting to avoid recovery of the floating fat layer. A portion of contaminating fat was extracted by passing the supernatant through glass wool loosely packed in a Pasteur pipette. The supernatant from the 30,000 x g spin was further clarified by centrifugation at 100,000 x g for 60 min. All centrifugation steps were performed at 5°C. The resulting supernatant was used immediately for desaturation assays described below or frozen in aliquots in liquid N₂ and stored at -70°C until further use. Of note, extracts developed a brown color, presumably due to extensive phenolic oxidation, when maintained at -20°C for longer than 1-2 weeks.

Acyl-ACP desaturation assays were based on those previously described (8, 9). Assays were performed in a total volume of 150 µl in loosely capped 13 x 100-mm glass tubes and consisted of 1.25 mM NADPH (from a freshly prepared stock in 100 mM Tricine, pH 8.2), 3.3 mM ascorbate, 0.7 mM dithiothreitol, 8000 units of bovine liver catalase (Sigma), 5 µg of bovine serum albumin (Fraction V) (Sigma), 20 µg of spinach ferredoxin (Sigma), 80 milliunits of spinach ferredoxin:NADPH reductase (Sigma), 33 mM PIPES, pH 6.0, and 118 pmol of [1-¹⁴C] acyl-ACP or -CoA. Reactions were started with the addition of the 100,000 x g supernatant of homogenized *T. alata* endosperm (typically 20-25 µg of total protein) and were conducted at room temperature (-22°C) with shaking (100 revolutions/min). Assays were terminated with the addition of 850 µl of 2.35 M NaOH and carrier fatty acids (30 µg of palmitic and petroselinic acid). The stopped reactions were then heated at 85°C for 1 hour. Following acidification with 350 µl of 4M H₂SO₄, the resulting fatty acids were recovered by three extractions with 2.5 ml

of hexane. Fatty acids were converted to methyl ester derivatives with 10% (w/v) boron trichloride in methanol (Alltech) using the method described by Morrison and Smith (24). Reaction products were then analyzed on 15% AgNO₃ TLC plates developed sequentially to heights of 10 and 20 cm in toluene at -20°C. TLC plates were prepared as described previously (25). Radioactivity was detected by autoradiography and quantified by liquid scintillation counting of TLC scrapings in a non-aqueous complete mixture.

To confirm the identity of 16:1Δ⁶ produced from palmitoyl-ACP, assays were conducted as described above using [U-¹⁴C] palmitoyl-ACP as the substrate. The methyl ester derivative of the monounsaturated product was purified by argentation TLC as described above and eluted from TLC scrapings with hexane/ethyl ether (2:1, v/v). The monounsaturated methyl ester was then cleaved at its double bond using permanganate-periodate oxidation (26). Chain lengths of oxidation products were determined relative to [¹⁴C] fatty acid standards by reverse-phase TLC using a mobile phase of acetonitrile/methanol/water (75:25:0.5).

Inhibition of desaturase activity was examined by supplementing assays with 1 mM KCN (neutralized) or 1 mM H₂O₂. In the latter case, catalase was omitted from reactions. Oxygen dependence of desaturase activity was characterized by purging assay tubes completely with nitrogen prior to and after addition of plant extract, and the reaction tube was tightly capped for the duration of the assay.

Radiolabeled acyl-ACPs were synthesized enzymatically using *Escherichia coli* ACP according to the method of Rock and Garwin (27). The following fatty acids were used in the synthesis of acyl-ACPs: [1-¹⁴C] myristic acid (American Radiolabeled Chemicals, St. Louis, MO) (specific activity 55 mCi/mmol), [1-¹⁴C] palmitic acid (NEN Dupont) (specific activity 58 mCi/mmol), [U-¹⁴C] palmitic acid (NEN Dupont) (specific activity 800 mCi/mmol), and [1-¹⁴C] stearic acid (American Radiolabeled Chemicals) (specific activity 55 mCi/mmol). [1-¹⁴C] Palmitoyl-CoA (specific activity 52 mCi/mmol) was purchased from Amersham Corp. A [¹⁴C] petroselinic acid

standard was prepared by incubation of conifer endosperm slices in [$1\text{-}^{14}\text{C}$] acetate as described previously (25).

***T. alata* Endosperm cDNA Library Construction** - Total RNA was isolated from *T. alata* endosperm using the method of Hall et al. (28). RNA was then passed through a cellulose (Sigma Cell 50, Sigma) column in order to reduce amounts of polysaccharides potentially recovered along with the RNA.

Poly(A)⁺ RNA was enriched by passing total RNA once through a column of oligo (dT) cellulose (Pharmacia LKB Biotechnology Inc.) and subsequently used in the construction of a Uni-ZAP XR (Stratagene) cDNA expression library according to the instructions of the manufacturer. A portion of the total amplified library packaged in phage was mass excised (29) yielding pBluescript II SK(-) harboring cDNA inserts. The recovered plasmid DNA was used for cDNA isolation by colony hybridization and polymerase chain reaction (PCR) amplification as described below.

PCR Amplification of Nucleotide Sequences Encoding Acyl-ACP Desaturases - Fully degenerate sense and antisense oligonucleotides were prepared that corresponded respectively to the conserved amino acid sequences Gly-Asp-Met-Ile-Thr-Glu-Glu and Glu-Lys-Thr-Ile-Gln-Tyr-Leu present in Δ^9 18:0 - (13, 15-20) and Δ^4 16:0-ACP desaturases (21). The sequence of the resulting sense and antisense oligonucleotides were 5'-GG(A/C/G/T)GA(C/T)ATGAT(A/C/T)AC(A/C/G/T)GA(A/G)GA-3' and 5'-A(A/G)(A/G)TATTG(A/G/T)AT(A/C/G/T)GT(C/T)TT(C/T)TC-3', respectively. Included on the 5' terminus of each oligonucleotide was sequence (5'-CAUCAUCAUCAU-3' or 5'-CUACUACUACUA-3') that allowed for insertion of PCR products into the pAMP1 vector (Life Technologies, Inc.). Template for PCR amplification was generated by transformation of the SOLR strain (Stratagene) of *E. coli* with an aliquot of the mass-excised *T. alata* endosperm cDNA library. Following growth of transformed *E. coli* to stationary phase in 3 ml of liquid culture, plasmid DNA was purified for use as template in PCR amplification. Reactions were performed in a 50- μl volume and consisted of 10 μM sense and antisense oligonucleotides, 150-300 ng of plasmid DNA

derived from the *T. alata* cDNA library. 2 mM MgCl₂, 0.2 mM dNTPs, 1 x *Taq* reaction buffer (Life Technologies, Inc.), and 5 units of *Taq* polymerase (Life Technologies, Inc.). Temperature conditions for PCR amplification were 5 min at 95°C and 25 cycles of 1 min at 95°C, 1.5 min at 55°C, and 1.5 min at 72°C. This was followed by an additional 10 min extension at 72°C. PCR fragments of approximately 215 base pairs were gel-purified, ligated into the pAMP1 vector using the CloneAmp system (Life Technologies, Inc.) according to the manufacturer's protocol, and introduced into *E. coli* DH5a (Life Technologies, Inc.). The resulting colonies were screened using colony hybridization as described by Sambrook et al. (30). A "negative" screening protocol was used to reduce the chances of reisolating cDNAs (pTAD1, 2, and 3) encoding Δ^8 18:0-ACP desaturases that were previously obtained by antibody screening of the *T. alata* endosperm cDNA library (19). DNA probes for library screening were formed by PCR amplification of portions of pTAD 1, 2, and 3. Primers and PCR reaction conditions were the same as those described above. An equimolar mixture of the PCR products derived from pTAD1, 2, and 3 was used as template for the synthesis of [α -³²P]dCTP random-primed labeled probes. Hybridization of plasmids of lysed colonies with radiolabeled probes was carried out in 6 x SSC and 0.25% (w/v) non-fat dry milk with shaking for 4 hours at 53°C as described by Sambrook et al. (30). Filters were washed three times in 1 x SSC and 0.1% SDS at 60°C (45 min/wash) and exposed to autoradiography. Plasmid DNA was subsequently isolated from 10 colonies which displayed little or no hybridization to the probes. Nucleotide sequence of the inserts of these plasmids was obtained by dideoxy chain termination using Sequenase 2.0 (United States Biochemical Inc.) according to the manufacturer's instructions. Two classes of plasmids were identified (designated pEC6 and 7), both of which contained inserts encoding portions of apparent acyl-ACP desaturases (based on amino acid identity with known Δ^8 18:0- and Δ^4 16:0-ACP desaturases).

Screening of a *T. alata* Endosperm cDNA Library for a Full-length Divergent Acyl-ACP Desaturase - Aliquots of the mass excised *T. alata* endosperm cDNA library were used to transform *E. coli* SOLR cells.

Approximately 50,000 colonies were screened using colony hybridization as described previously (30). Nucleotide probes for screening were generated by [α - 32 P]dCTP random-primed hexamer labeling of inserts of pEC6 and 7. Hybridization and washing conditions were the same as those described above. Colonies containing plasmid DNA that strongly hybridized to the probe derived from pEC6 were isolated, and nucleotide sequence was obtained from both strands of the longest cDNA insert (the corresponding plasmid was designated pTAD4) using Sequenase 2.0. Because of a relative lack of abundance, colonies containing plasmid hybridizing to the pEC7-derived probe were not further characterized.

***E. coli* Expression of a Putative cDNA for Δ^6 Palmitoyl-ACP Desaturase** - To determine the activity of the desaturase encoded by pTAD4, the portion of the clone corresponding to the mature peptide (total protein minus plastid transit peptide) was expressed in *E. coli*. This region of the cDNA insert of pTAD4 was first amplified by PCR using Vent DNA polymerase (New England Biolabs). The nucleotide sequence of the sense primer was 5'-GCTTCGACTATTACTCAC3'. M-13(-20) forward primer was used as the antisense oligonucleotide. The PCR product was blunt-end ligated into the *Nco*I site of the *E. coli* expression vector pET3d (Novagen) as described (30). The *Nco*I-digested vector had been previously treated with the Klenow fragment of DNA polymerase I to fill-in 5' protruding ends. The junction between the vector and the 5' terminus of the insert was sequenced to confirm that the PCR product was ligated into pET3d in the proper reading frame. This construct was subsequently introduced into the *E. coli* strain BL21 pLysS and grown in LB media with carbenicillin (125 μ g/ml) and chloroamphenicol (30 μ g/ml) selection. At a cell density of OD₆₀₀ ~ 0.8, cultures were induced with the addition of isopropyl-1-thio- β -D-galactopyranoside to a final concentration of 0.5 mM and grown for an additional 4 hours. Cells were then washed in 50 mM Tris-HCl, pH 7.5, lysed by two freeze-thaw cycles (using a liquid nitrogen bath for freezing and a 22°C water bath for thawing). Lysates were then incubated with bovine pancreas DNase I (Boehringer Mannheim) (20 μ g/ml) for 15 min at 22°C. The

extract was subsequently centrifuged at 14,000 x g for 5 min. The resulting supernatant was used for acyl-ACP desaturation assays as described above. Radiolabel in the TLC-analyzed reaction products was detected using a Bioscan System 200 image scanner. The double bond position of the monounsaturated product was determined by gas chromatography-mass spectrometry analysis of its dimethyl disulfide derivative (31). In these studies, the desaturation assays described above were scaled up 6-fold, and reactions were conducted with 2.6 nmol of [1-¹⁴C] 16:0-ACP and 1.1 mg of soluble protein of lysed *E. coli* expressing the *T. alata* cDNA. Assays were conducted for 4 hours. High protein concentrations and long incubation periods were used to ensure the synthesis of sufficient amounts of monounsaturated fatty acid for mass spectral analyses. Reaction products were converted to fatty acid methyl esters as described above and subsequently reacted with 100 μ l of an iodine solution (60 mg/ml ethyl ether) and 350 μ l of dimethyl disulfide (Aldrich). After 3 hours of incubation with shaking (250 revolutions/min) at 37°C, dimethyl disulfide derivatives of unsaturated fatty acid methyl esters were extracted as described previously (32). These derivatives (dissolved in hexane) were then analyzed by gas chromatography-mass spectrometry using a Hewlett Packard PH5890II gas chromatograph interfaced with a HP5971 mass selective detector. Separation of analytes was achieved using a DB23 (30 m x 0.25 mm inner diameter) column (J&W Scientific) with the oven temperature programmed from 185°C (3 min hold) to 230°C at rate of 2.5°C/min.

Results

Detection of a Soluble Δ^6 Palmitoyl-ACP Desaturase in *T. alata* Endosperm Extracts - The seed oil of *T. alata* is composed of nearly 85% weight of the unusual monounsaturated fatty acid Δ^6 hexadecenoic acid (16:1 Δ^6). To examine the metabolic origin of the double bond of this fatty acid, the 100,000 x g supernatant of a homogenate of developing *T. alata* seed endosperm was incubated with [¹⁴C]16:0-ACP and potential desaturase cofactors. Using this assay system, substantial amounts of 16:0-ACP desaturase activity were detected in the soluble endosperm extract (Fig. 1).

In the absence of a radiolabeled standard for 16:1 Δ^6 , two independent analytical methods indicated that the double bond of the resulting 16:1 moiety was positioned at the Δ^6 carbon atom: 1) the 16:1 desaturation product displayed mobility on argentation TLC plates similar to that of the Δ^6 monounsaturated fatty acid petroselinic acid (18:1 Δ^6) when these molecules were analyzed as methyl ester derivatives (Fig. 1) and 2) permanganate-periodate oxidation of the methyl ester of the 16:1 desaturation product gave rise to a molecule with mobility on reverse-phase TLC equivalent to that of decanoic acid (10:0) (Fig. 2, *Product B*) as well as to an acyl moiety containing a lesser number of carbon atoms (Fig. 2, *Product A*).

Substrate Properties of the Δ^6 Acyl-ACP Desaturase - To confirm that the Δ^6 desaturase identified above is most active with 16:0-ACP, assays were conducted using ^{14}C -saturated acyl-ACP substrates containing 14, 16, and 18 carbons. As with 16:0-ACP (described above), Δ^6 desaturase activity was also detected when [1- ^{14}C]14:0- and 18:0-ACP were reacted with a 100,000 x g supernatant of a *T. alata* endosperm homogenate. Following derivitization, a portion of the desaturation products resulting from 18:0-ACP comigrated on argentation TLC with the methyl ester of petroselinic acid, which was resolvable in this system from methyl oleic acid (data now shown). Similarly, the desaturation product arising from 14:0-ACP migrated in the expected position for a fatty acid containing a Δ^6 double bond (data not shown). Therefore, it appears that the Δ^6 desaturase of *T. alata* endosperm positions the placement of unsaturation with regard to the carboxyl end of fatty acid substrates. This double bond positioning property has been previously observed with the Δ^9 18:0- and Δ^4 16:0-ACP desaturases (22, 33).

Under the assay conditions used, Δ^6 16:0-ACP desaturase activity in the 100,000 x g supernatant of a *T. alata* endosperm homogenate was essentially linear over 10 min (Fig. 3). When assays were conducted over this time period, the specific activity of the Δ^6 desaturase was approximately 7-fold higher using [1- ^{14}C]16:0-ACP as a substrate rather than either [1- ^{14}C]14:0- or 18:0-ACP (Table 1). Values obtained with the latter substrate, however, were obscured because of the presence of competing Δ^9 18:0-ACP

desaturase activity in the endosperm extract. Finally, no desaturase activity was detected when $[1-^{14}\text{C}]16:0\text{-CoA}$ was presented as a potential substrate. Overall, these results indicate that the Δ^6 desaturase is most active *in vitro* with 16:0 esterified to ACP.

TABLE I

In vitro substrate specificities of acyl-ACP or -CoA desaturases of
T. alata endosperm

Desaturase assays were conducted for 10 min using 118 pmol of $[1-^{14}\text{C}]$ acyl-ACP or -CoA substrate and 23 μg of total protein from 100,000 \times g supernatant of a *T. alata* endosperm homogenate.

<u>Substrate</u>	<u>Monounsaturated products^a</u>	
	Δ^6	Δ^9
	<i>pmol/min/mg protein</i>	
14:0-ACP	13	ND ^b
16:0-ACP	99	ND
16:0-CoA	ND	ND
18:0-ACP	12	173 ^c

^a14:1 Δ^6 , 18:1 Δ^6 , and 18:1 Δ^9 were identified by the mobilities of these fatty acids on argention TLC plates.

^bNot detected.

^cAssay conditions were adjusted only for the linear measurement of Δ^6 desaturase activity. Therefore this value may underestimate the specific activity of Δ^9 18:0-ACP desaturase.

Cofactors and Inhibitors of Δ^6 Palmitoyl-ACP Desaturase Activity -

Additional *in vitro* assays were conducted to compare the functional properties of the Δ^6 16:0-ACP desaturase with those previously determined for the Δ^9 18:0-ACP desaturase (7, 8, 10). In this regard, virtually no Δ^6 16:0-ACP desaturase activity was detected in the 100,000 x g supernatant of *T. alata* endosperm homogenates when assays were conducted in the absence of ferredoxin or molecular oxygen (Fig. 4). Δ^6 16:0-ACP desaturase activity was also reduced when catalase was omitted from assays. Furthermore, the inclusion of 1 mM KCN or H_2O_2 in reactions resulted in the loss of most of the desaturase activity. Such catalytic properties of the *T. alata* Δ^6 16:0-ACP desaturase were similar to those previously described for the Δ^9 18:0-ACP desaturase (7, 8, 10).

Isolation of a cDNA Encoding a Diverged Acyl-ACP Desaturase from *T. alata* Endosperm - Based on functional similarities of the Δ^6 16:0- and Δ^9 18:0-ACP desaturases described above, we examined whether these enzymes are also structurally related. To address this question, attempts were made to isolate a cDNA for the Δ^6 16:0-ACP desaturase using Δ^9 18:0-ACP desaturase-derived probes. As a first approach, a cDNA expression library prepared from poly(A)⁺ RNA of *T. alata* endosperm was screened with antibodies against the Δ^9 18:0-ACP desaturase of avocado (13). This method was previously used to obtain a cDNA for the Δ^4 16:0-ACP desaturase of coriander endosperm (21). In the present study, however, antibody screening of the *T. alata* endosperm expression library yielded only cDNAs for three apparent isoforms of the Δ^9 18:0-ACP desaturase, which were designated pTAD1, 2, and 3 (19).

As an alternative approach, PCR amplification of a Δ^6 16:0-ACP desaturase-specific nucleotide probe was attempted using degenerate sense and antisense oligonucleotides prepared against two conserved amino acid sequences in Δ^9 18:0- and Δ^4 16:0-ACP desaturases. One of the sequences (Gly-Asp-Met-Ile-Thr-Glu-Glu) is encoded by the cDNA for the Δ^6 16:0-ACP desaturase and all known cDNAs for the Δ^9 18:0-ACP desaturase. The second sequence (Glu-Lys-Thr-Ile-Gln-Tyr-Leu) is also encoded by the

Δ^4 16:0-ACP desaturase of cDNA and all known Δ^9 18:0-ACP desaturase cDNAs except that of safflower (14). Products of approximately 215 base pairs obtained following one round of PCR amplification of the total *T. alata* cDNA library (in plasmid form) were screened after subcloning into the pAMP1 vector. To delineate products of the previously isolated cDNAs pTAD1, 2, and 3, colonies containing PCR-derived clones were screened in a negative manner with random-labeled probes for pTAD1, 2, and 3 and conditions of moderate to high stringency. One of the resulting clones (pEC6) that displayed weak or no hybridization to these probes encoded an amino acid sequence that was somewhat diverged from those of known Δ^9 18:0-ACP desaturases.

When the *T. alata* endosperm library was screened with a random-labeled probe prepared from the insert of pEC6, >0.1% of the total cDNAs examined strongly hybridized to this probe. The longest of a selected portion of these cDNAs (the corresponding plasmid was designated pTAD4) contained 1279 base pairs and had an open-reading frame corresponding to a 387-amino-acid polypeptide with considerable identity to known Δ^4 16:0- and Δ^9 18:0-ACP desaturases (Fig. 5). Based on similarity of flanking bases to the consensus sequence proposed by Lütcke et al. (34), the translational start site of the cDNA insert of pTAD4 likely occurs at nucleotide 17. In addition, from homology with Δ^4 16:0- and Δ^9 18:0-ACP desaturases, the mature peptide encoded by pTAD4 likely begins at amino acid 33. As such, the 32 amino acids preceding this residue correspond to a putative plastid transit peptide as is present in all acyl-ACP desaturases characterized to date.

Interestingly, the cDNA insert of pTAD4 lacks nucleotide sequence for 6-7 amino acids found near the amino terminus of all previously characterized Δ^9 18:0-ACP desaturases. This region is also altered in the cDNA for the coriander Δ^4 16:0-ACP desaturase (21) as compared to cDNAs for Δ^9 18:0-ACP desaturases. In this case, the coding sequence for 15 amino acids is absent in the Δ^4 16:0-ACP desaturase cDNA relative to the castor Δ^9 18:0-ACP desaturase cDNA (13) (Fig. 5). The pTAD4-encoded peptide also contains 2 less amino acids at its carboxyl terminus than both the Δ^4 16:0- and Δ^9 18:0-

ACP desaturases. Despite these differences, the interior regions of the putative desaturase encoded by pTAD4 share significant identity with portions of the primary structures of Δ^4 16:0- and Δ^9 18:0-ACP desaturases, and the spacing between conserved regions of amino acids is the same in all three desaturase types. Overall, the mature peptide encoded by the cDNA insert of pTAD4 shares 66% identity with the castor Δ^9 18:0-ACP desaturase and 57% identity with the coriander Δ^4 16:0-ACP desaturase, disregarding any missing amino acids.

Activity of an *E. coli*-expressed cDNA for a Diverged Acyl-ACP Desaturase of *T. alata* Endosperm - To determine the activity of the desaturase corresponding to the cDNA insert of pTAD4, the mature peptide-encoding region of this clone was expressed in *E. coli* with expression driven by the T7 RNA polymerase promoter of the vector pET3d (Novagen). When assayed with [$1\text{-}^{14}\text{C}$]16:0-ACP, crude extracts of isopropyl-1-thio- β -D-galactopyranoside-induced recombinant *E. coli* catalyzed the synthesis of [^{14}C]16:1 (Fig. 6). In addition, the methyl ester of the 16:1 product displayed mobility on argentation TLC plates similar to that of a methyl petroselinic acid (18:1 Δ^6) standard, suggesting that this monounsaturated product is a Δ^6 isomer. Detectable acyl-ACP desaturase activity was absent in extracts of *E. coli* harboring the pET3d vector without cDNA insert or in uninduced recombinant *E. coli*. Furthermore, like the activity found in *T. alata* endosperm extracts, the desaturase expressed in *E. coli* displayed an *in vitro* substrate preference for 16:0-ACP and exhibited no detectable activity in the absence of reduced ferredoxin (Fig. 7).

The [^{14}C]16:0 moiety produced *in vitro* from the *E. coli*-expressed desaturase was conclusively identified as a Δ^6 isomer through gas chromatography-mass spectrometry analysis of its dimethyl disulfide derivative (Fig. 8). In the mass spectrum shown, the ions 145, 177, 187, and 364 *m/z* are diagnostic for a [$1\text{-}^{14}\text{C}$]16:1 Δ^6 moiety. Significant amounts of non-radiolabeled or [^{12}C]16:1 Δ^6 were also detected among the desaturase assay products. This was indicated by the presence of the additional ions 143, 175, and 362 *m/z* in the mass spectrum of [$1\text{-}^{14}\text{C}$]16:1 Δ^6 as well as by an

enrichment in the abundance of ions 187 m/z (Fig. 8). It is unlikely that the non-radiolabeled 16:1 Δ^6 resulted from *in vivo* synthesis in *E. coli*. In this regard, *E. coli* does not normally produce 16:1 Δ^6 (35). Furthermore, gas chromatographic analysis of fatty acids of *E. coli* expressing the pTAD4-encoded desaturase failed to detect any 16:1 Δ^6 in the bacterial lipids (data now shown). Given the relatively high concentrations of *E. coli* protein used in these assays, unlabeled 16:1 Δ^6 likely arose from the *in vitro* desaturation of endogenous *E. coli* 16:0-ACP present in crude bacterial extracts.

Of note, expression levels of the *T. alata* cDNA in *E. coli* appeared to be low relative to that often obtained with DNA inserts placed behind the T7 RNA polymerase promoter (36). The expressed protein, for example, could not be distinguished on Coomassie-stained SDS-polyacrylamide gels of either the total soluble or insoluble protein fractions of lysed *E. coli* (data not shown). Also suggestive of low expression levels in *E. coli*, the specific activity of Δ^6 16:0-ACP desaturase in recombinant *E. coli* extracts (Fig. 7) was typically half of that detected in *T. alata* endosperm homogenates (Table I).

The results presented here demonstrate the involvement of a novel soluble Δ^6 16:0-ACP desaturase in the synthesis of Δ^6 hexadecenoic acid in the endosperm of *T. alata*. The activity of this enzyme has several properties similar to those previously described for the Δ^9 18:0-ACP desaturase. These include the requirement of reduced ferredoxin for detectable *in vitro* activity, the stimulation of activity by catalase, and the inhibition of activity by potassium cyanide and hydrogen peroxide. The existence of a Δ^6 16:0-ACP desaturase in *T. alata* endosperm was confirmed by the isolation of a cDNA for this enzyme. While the amino acid sequence deduced from this cDNA shares some identity with Δ^9 18:0- and Δ^4 16:0-ACP desaturases, these findings, together with those previously obtained for petroselinic acid biosynthesis (19, 21), indicate that natural variations in the primary structures of acyl-ACP desaturases can give rise to novel enzymes with altered substrate recognition and double bond positioning properties.

The major difference between the primary structures of the mature Δ^6 16:0-, Δ^4 16:0-, and Δ^9 18:0-ACP desaturases occurs in a region near their

amino termini. In this region, the *T. alata* Δ^9 16:0-ACP desaturase contains 6 less amino acids than the castor Δ^9 18:0-ACP desaturase. Similarly, this portion of the coriander Δ^9 16:0-ACP desaturase lacks 15 amino acids relative to the castor Δ^9 18:0-ACP desaturase. Without intending to be limited to any particular theory, one possibility is that differences in recognition of substrate chain length (16:0-ACP *versus* 18:0-ACP) and/or double bond positioning of these desaturases are associated with this divergence in the primary structures of these enzymes. Alternatively, this region of the amino terminus of Δ^9 18:0-ACP desaturase may not contribute significantly to the catalytic properties of this enzyme. As such, while again not intending to be limited to any particular theory, if the Δ^4 and Δ^6 16:0-ACP desaturases evolved from the Δ^9 18:0-ACP desaturase, then there may have been little selective pressure to maintain this region intact in the variant 16:0-ACP desaturases.

Ultimately, an understanding of how differences in the amino acid sequences of Δ^4 16:0-, Δ^6 16:0-, and Δ^9 18:0-ACP desaturases contribute to variations in their functional properties will require comparisons of the three-dimensional structures of these enzymes. In this regard, elucidation of the crystal structure of the castor Δ^9 18:0-ACP desaturase is currently in progress (37). With such information, it will be possible to overlap amino acid sequences of the Δ^4 and Δ^6 16:0-ACP desaturases onto the three-dimensional structure of Δ^9 18:0-ACP desaturase to more precisely identify residues associated with the different substrate recognition and double bond positioning properties of these enzymes. This could eventually lead to the design of "tailor-made" desaturases that are capable of inserting double bonds into a variety of positions of acyl moieties of a range of carbon chain lengths.

An interesting observation from the studies described above was the lack of detectable amounts of 16:1 Δ^6 in lipids of *E. coli* expressing the *T. alata* cDNA. Similarly, Thompson et al. (14) reported that expression of the safflower Δ^9 18:0-ACP desaturase cDNA did not lead to the *in vivo* production of oleic acid in recombinant *E. coli*. The latter result can be explained by the fact that *E. coli* contains little 18:0-ACP (38). However, 16:0-ACP is a major

component of the acyl-ACP pool of *E. coli*. Therefore it is unlikely that the lack of 16:1 Δ^6 synthesis in *E. coli* expressing the *T. alata* cDNA is due to the presence of insufficient substrate for the desaturase. In addition, *E. coli* has been reported to contain ferredoxin (39), the apparent electron donor for the Δ^6 16:0-ACP desaturase. However, as proposed by Thompson et al. (14), *E. coli* ferredoxin may not functionally interact with plant acyl-ACP desaturases. Alternatively, *E. coli* may not have adequate amounts of ferredoxin in a reduced form as required for Δ^6 16:0-ACP desaturase activity.

In addition to 16:1 Δ^6 , *T. alata* seed contains the unusual fatty acid 18:1 Δ^8 , which composes about 2% weight of the oil of this tissue (23). We have previously shown that petroselinic acid (18:1 Δ^6) is formed by elongation of 16:1 Δ^4 -ACP in Umbelliferae endosperm (22). In an analogous manner, we predict that 18:1 Δ^8 arises from the elongation of 16:1 Δ^6 -ACP rather than from the Δ^8 desaturation of 18:0-ACP. Unlike the synthesis of petroselinic acid, though, elongation of 16:1-ACP in *T. alata* endosperm is likely not a major pathway as the ratio of amounts of 16:1 Δ^6 :18:1 Δ^8 in this tissue is approximately 40:1. In contrast, the ratio of amounts of 16:1 Δ^6 :18:1 Δ^6 in endosperm of the Umbelliferae coriander is more than 1:500 (22, 25).

Finally, significant efforts have been directed toward the development of transgenic crops that produce high value specialty oils (4, 40, 41). Using methodologies currently well known in the art, transgenic plants could be produced (42, 43) which would contain and express the Δ^6 16:0-ACP desaturase gene and which would produce high levels of 16:1 Δ^6 . In this regard, oils rich in 16:1 Δ^6 may have properties suitable for industrial use. Like petroselinic acid, 16:1 Δ^6 can be oxidatively cleaved at its double bond to yield adipic acid, a precursor of nylon 6,6. In addition, high palmitic acid (16:0) mutants of crop plants including soybean (44) and *Brassica campestris* (45) are available that could serve as appropriate backgrounds for transgenic expression of the cDNA for the *T. alata* Δ^6 16:0-ACP desaturase. Still the success of such research would likely require additional studies to determine whether enzymes other than Δ^6 16:0-ACP desaturase are

specialized for the synthesis and metabolism of 16:1 Δ^5 in *T. alata* endosperm. For example, a petroselinoyl-ACP-specific thioesterase has been identified in Umbelliferae endosperm extracts that efficiently releases petroselinic acid from ACP and, as a result, makes this fatty acid available for subsequent storage in triacylglycerol (46). A related enzyme may also be required for high levels of 16:1 Δ^6 accumulation in transgenic plants.

R e f e r e n c e s

1. Cheesbrough, T. M. (1990) *Plant Physiol.* **93**, 555-559.
2. Wada, H., Gombos, Z., and Murata, N. (1990) *Nature* **347**, 200-203
3. Miquel, M., James Jr., Dr. Dooner, H., and Browse, J. (1993) *Proc. Natl. Acad. Sci. U.S.A.* **90**, 6208-6212.
4. Battey, J. F., Schmid, K. M., and Ohlrogge, J. B. (1989) *Trends Biotech.* **7**, 122-125.
5. Heinz, E. (1993) in *Lipid Metabolism in Plants* (Moore, T. S., ed) pp. 33-89, CRC Press, Boca Raton, FL.
6. Nagai, J., and Bloch, K. (1965) *J. Biol. Chem.* **240**, 3702-3703.
7. Nagai, J., and Bloch, K. (1968) *J. Biol. Chem.* **243**, 4626-4633.
8. Jaworski, J. G., and Stumpf, P. K. (1974) *Arch. Biochem. Biophys.* **162**, 158-165.
9. McKeon, T. A., and Stumpf, P. K. (1981) *Methods Enzymol.* **71**, 275-281.
10. McKeon, T. A., and Stumpf, P. K. (1982) *J. Biol. Chem.* **257**, 12141-12147.
11. Cheesbrough, T. M., and Cho, S. H. (1990) in *Plant Lipid Biochemistry: Structure and Utilization* (Quinn, P. J., and Harwood, J. L., eds) pp. 129-130, Portland Press, London.
12. Kinney, A. J., Hitz, W. D., and Yadav, N. S. (1990) in *Plant Lipid Biochemistry: Structure and Utilization* (Quinn, P. J., and Harwood, J. L., eds) pp. 126-128, Portland Press, London.
13. Shanklin, J., and Somerville, C. (1991) *Proc. Natl. Acad. Sci. U.S.A.* **88**, 2510-2514.
14. Thompson, G. A., Scherer, D. E., Foxall-Van Aken, S., Kenny, J. W., Young, H. L., Shintani, D. K. Kridl, J. C., and Knauf, V. C. (1991) *Proc. Natl. Acad. Sci. U.S.A.* **88**, 2578-2582.
15. Nishinda, I., Beppu, T., Matsuo, T., and Murata, N. (1992) *Plant Mol. Biol.* **19**, 711-713.

16. Sato, A., Becker, C. K. and Kanuf, V. C. (1992) *Plant Physiol.* **99**, 362-363.
17. Slocombe, S. P., Cummins, I., Jarvis, R. P., and Murphy, D. J. (1992) *Plant Mol. Biol.* **20**, 151-155.
18. Taylor, M. A., Smith, S. B., Davies, H. V., and Burch, L. R. (1992) *Plant Physiol.* **100**, 533-534.
19. Cahoon, E. B., Becker, C. K. Shanklin, J., and Ohlrogge, J. B. (1994) *Plant Physiol.* in press.
20. Singh, S., McKinney, S., Green, A. (1994) *Plant Physiol.* **104**, 1075.
21. Cahoon, E. B., Shanklin, J., and Ohlrogge, J. B. (1992) *Proc. Natl. Acad. Sci. U.S.A.* **89**, 11184-11188.
22. Cahoon, E. B., and Ohlrogge, J. B. (1994) *Plant Physiol.* **104**, 827-838.
23. Spencer, G. F., Kleiman, R., Miller, R. W., and Earle, F. R. (1971) *Lipids* **6**, 712-714.
24. Morrison, W. R. and Smith, L. M. (1964) *J. Lipid Res.* **5**, 600-608.
25. Cahoon, E. B., and Ohlrogge, J. B. (1994) *Plant Physiol.* **104**, 845-855.
26. Christie, W. W. (1982) *Lipid Analysis*, 2nd Ed., Pergammon Press, Oxford.
27. Rock, C. O., and Garwin, J. L. (1979) *J. Biol. Chem.* **254**, 7123-7128.
28. Hall, T. C., Ma, Y., Buchbinder, B. U., Pyne, J. W., Sun, S. M., and Bliss, F. A. (1978) *Proc. Natl. Acad. Sci. U.S.A.* **75**, 3196-3200.
29. Hay, B., and Short, J. M. (1992) *Strategies* **5**, 16-18.
30. Sambrook, J., Fritsch, E. F., and Maniatis, T. (1989) *Molecular Cloning: a Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
31. Francis, G. W. (1981) *Chem. Phys. Lipids* **29**, 369-374.
32. Yamamoto, K. Shibahara, A., Nakayama, T., and Kajimoto, G. (1991) *Chem. Phys. Lipids* **60**, 39-50.
33. Gibson, K. J. (1993) *Biochim. Biophys. Acta* **1169**, 231-235.
34. Lütcke, H. A., Chow, K. C., Mickel, F. S., Moss, K. A., Kern, H. F., and Scheele, G. A. (1987) *EMBO J.* **6**, 43-48.

35. Cronan Jr., J. E., and Rock, C. O. (1987) in *Escherichia coli and Salmonella typhimurium Cellular and Molecular Biology* (Neihardt, F. C., ed) pp. 474-497. American Society of Microbiology, Washington, D.C.
36. Studier, F. W., Rosenberg, A. H., Dunn, J. J., and Dubendorff, J. W. (1990) *Methods Enzymol.* **185**, 60-89.
37. Schneider, G., Lindqvist, Y., Shanklin, J., and Somerville, C. (1992) *J. Mol. Biol.* **225**, 561-564.
38. Rock, C. O., and Jackowski, S. (1982) *J. Biol. Chem.* **257**, 10579-10765.
39. Knoell, H. E., and Knappe, J. (1974) *Eur. J. Biochem.* **50**, 245-252.
40. Murphy, D. J. (1992) *Trends Biotechnol.* **10**, 84-87.
41. Ohlrogge, J. B. (1992) *Plant Physiol.* **104**, 821-826.
42. Schilperoort, R. A., Hoekema, A., and Hooykaas, P. J. J. (1990) U. S. Patent No. 4,940,838.
43. Tomes, D., Bidney, D., Buising C. M. (1994) U. S. Patent No. 5,322,783.
44. Bubeck, D. M., Fehr, W. R. and Hammond, E. G. (1980) *Crop. Sci.* **29**, 652-656.
45. Perrson, C. (1985) *Cruciferae Newsletter* **10**, p. 137, Scottish Crop Research Institute, Invergowrie, Dundee, Scotland.
46. Dörmann, P., Frentzen, M., and Ohlrogge, J. B. (1994) *Plant Physiol.* **104**, 839-844.

What Is Claimed Is:

1. An isolated Δ^6 - palmitoyl-acyl carrier protein desaturase.
2. A transgenic plant cell comprising heterologous DNA encoding a Δ^6 - palmitoyl-acyl carrier protein desaturase.
3. The transgenic plant cell of Claim 2 wherein said heterologous DNA is as shown in SEQ. I.D. NO.: 1.
4. The transgenic plant cell of Claim 2 wherein said Δ^6 - palmitoyl-acyl carrier protein desaturase is expressed and active.
5. The transgenic plant cell of Claim 2 as a seed or a propagate of the seed.
6. An isolated DNA encoding a Δ^6 - palmitoyl-acyl carrier protein desaturase.
7. The DNA of Claim 6 wherein said DNA encodes a *Thunbergia alata* Δ^6 - palmitoyl-acyl carrier protein desaturase.
8. The DNA of Claim 6 wherein said DNA is as shown in SEQ. ID NO.: 1.
9. A mRNA derived from a DNA as shown in SEQ. I.D. NO.: 1.
10. A transgenic plant comprising heterologous DNA encoding a Δ^6 - palmitoyl-acyl carrier protein desaturase.
11. The transgenic plant of Claim 10 wherein said heterologous DNA is as shown in SEQ. I.D. NO.: 1.
12. The transgenic plant of Claim 10 wherein said Δ^6 - palmitoyl-acyl carrier protein desaturase is expressed and active.
13. A vector comprising DNA encoding a Δ^6 - palmitoyl-acyl carrier protein desaturase.
14. A method for modifying the fatty acid content of a plant seed, the method comprising transforming the plant which produces the seed with a heterologous DNA encoding a Δ^6 - palmitoyl-acyl carrier protein desaturase, and introducing a double bond at the sixth carbon atom from the carboxyl end of a 16 carbon saturated fatty acid in the plant seed through the activity of the expressed desaturase.
15. The method of Claim 14 wherein the double bond at the sixth carbon atom from the carboxyl end of the 16 carbon saturated fatty acid is

introduced by a Δ^3 - palmitoyl-acyl carrier protein desaturase from *Thunbergia alata*.

16. The method of Claim 14 further comprising oxidative cleavage of the monounsaturated fatty acid formed by the method of Claim 14.
17. A plant seed having a modified fatty acid content derived through the process of Claim 14.
18. A plant seed having a modified fatty acid content derived through the process of Claim 15.
19. A plant seed having a modified fatty acid content derived through the process of Claim 16.

1/8




FIGURE 1

2/8




FIGURE 2

3/8




FIGURE 3

4/8




FIGURE 4

5/8

46	ATTTTGTAAAGTGAATGGCTTGGTATCAAGAGTATAGGAGCCATAAGACTCTCTCTGTACTTTAAATTAGCTTACCAGCTTTGTACCAC	97
46	N A L V F K S I G A N K T P P C T L N L A S P A L Y N	27
49	N A L K L N P F L S O T Q C L P S F A L P P H A S T R S	28
44	N A N K E L A L M T L Q C P K R N H F T R I A P P O A G R V R	31
46	ACCAGAGTCACATGGCTTCGACTATTACTCACCCTCCGCACTCAAGAT.....AGAAATATGCTACTCGACCA...GTAAGG.....	178
46	T R V T N A S I I T H P P L K D - - R K I S S T E R - V R - - -	54
49	P K F Y I I I L K S S S K E V E N L C I P F H P P P I I N V Q V	61
44	S K V S I I I L N A S I L V F I K L K A G R P - - E I - - -	57
46	ACATATCGTGGCTCCAGAGAGGCTGAATCTCAATTCTATGCAGGGTGGTTGAAGACCATCTCCCTTCTTGAAGCGGTGGAGGATCG	277
46	T Y P A P E K A E I F H S N H Q M V E D I L P P L K P V E E S	87
49	I N S H P I O I I I K I L D N I A I E N I I V H I I I I K C	94
44	- - - - - D I L I I L E I I A R I N I I V N I I I I N I	82
46	TGCAGCGCAGGAGCTTCTCCGCGCTCCGCTTCTGATGGTTCCAGAGCATGGAAGAGCTTCGTAACCAAGCGCGCATCTCCCTGATGATAC	376
46	H Q R T D F L P D S T S D G F A E D V E E L R K R T A D L P D D Y	120
49	I I I Q I I I P A I I I D I I I R I I I E I A K E I I I I	127
44	I I I Q I Y I I I P I I I A I E D I I K I N I E I A K I I I I E I	115
46	TTAGTTCATTTGGTGGAGCAATGGTACCGAGGAAGCCCTTCCGAGCTTCAACATGCTTAACACAGAGATGTGATATACATGAGAGCGGCCCC	475
46	L V A L V G A R V T E E A L P T Y Q T H L M I T T D V I Y D E S G A	153
49	F I I I I D I	160
44	F I V I I I D I	148
46	AGCCCTGTGCTTGGCGCTTTGGACCGCGCTTGGACCGCTGAAGAGACAGGCGATGGTATTTGTCAACAGATCTCTCTCTTTCCGCGTGTGTC	574
46	S P V P V A V V U T R A W T A E E N R H G D I V N K T L Y L S G R V	186
49	I I I T S I	193
44	Q I T S I	181
46	GATATGAAGCAATGAAGAGCTATCAATACTTGAATGGCTCGGCGATGATCTGTGGCGACACACCGCTACGATATATCTACAGCATCGC	673
46	D N K Q I E K T I O Y L I G S G N D P G A D N M P Y L A Y I Y T S	219
49	I I R I	226
44	I I R M I	214
46	TATCAGAGAGGGCTCAGCGATCTCCCATGGAAGTCTGGCGCGCTAGCGAGGAGAGGAGAGATGAACCTGGTCAGATTTGTGGACAACTTCT	772
46	Y Q E R A T A I S N G S L G R L A R Q K G E N K L A O I G C T I S	252
49	F I I I I F I I I I I N T A I Q I K E H D I I I I I I I I I I I	259
44	F I I I I F I I I I I A N T A K I I I Q E T I D E N I I I V I I I I I	247
46	GGCATGAGAGCGCGCAGCGCGCTACTCGAAATCGTGGAGAGCTATTCGAGTTGGTTCGACAGAGGCGACATGTTGGCGTTGGCATACATGATG	871
46	A D E K R R E A A Y S K I V E K L F E L D P E G T H L A L A Y M N	285
49	I I	292
44	S I I I I I A T I	280
46	AGATGAAGATTGTAAAGCAGCTCTGTCGATGACGATGGAGAGATCCGACATGTTCAACATTTCTGCTGTGTCGCGAGCGATGGGATTTAC	970
46	K N K I V M P A R L N H N D G K D P D F R H F S A V S Q R L G I T	318
49	R K I	325
44	R K I	313
46	ACTCAGAGAGGATACCGACATCTCGAGCATATGATAGCGCGTGGGAGTGGATAGCTAGCTAGCGGGCTGAGCGGGAGGGCGCGAGCGCGCAGGAT	1069
46	T A K Y I I D I K L T H N I A R M G V D K L T G L S E E R R A Q D	351
49	I I I D I A I I I I F L V G I I K I I I I I I I I I I I I I I I	358
44	S I W D I C I I I I D F L V D K I I N I A I N I I I I I I I I I I I	346
46	TACGTGTGCGGGTTGCCGATGAGGTAGGAAGGTGGAGGAGCGCGCGCGCTGGCGGAGATATATCGCAT...GTCTCTTTAGCTGATCTTT	1165
46	T V C S I P N I I R K V E R A D A V A E H I S N V P F S M I F	383
49	I I I R I	390
44	I I I S I A A K I	379
46	GGAGAGAGATG.....TAGTCTCAGTCTCAGTCTCAGTCTCAGTGTGTGTGTCTATGATCAAGAAATAGTGCAATGCCACCTTATTCTC	1258
46	G R R V - - -	387
49	D I Q I K I - - -	396
44	N I Q I I I I - - -	385

FIGURE 5

6/8




FIGURE 6

7/8




FIGURE 7

8/8




FIGURE 8

(51) International Patent Classification 6 : C12N 15/33, 15/82, 9/02, A01H 5/00		A3		(50) International Publication Number: WO 96/13591 (43) International Publication Date: 9 May 1996 (09.05.96)	
(21) International Application Number: PCT/US95/13784 (22) International Filing Date: 25 October 1995 (25.10.95)		(30) Priority Data: 08/329,560 26 October 1994 (26.10.94) US		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TH, TM, TT, UA, UG, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, LS, MW, SD, SZ, UG).	
(71) Applicant: MICHIGAN STATE UNIVERSITY [US/US]; 238 Administration Building, East Lansing, MI 48824 (US). (72) Inventors: CAHOON, Edgar, B.; Building 463, Brookhaven National Laboratories, Upton, NY 11973 (US). OHLROGGE, John, B.; 2123 Lagoon Drive, Okemos, MI 48864 (US). (74) Agents: YATES, Michael, E. et al.; 700 Capital Square, 400 Locust Street, Des Moines, IA 50309 (US).		Published With international search report.		(88) Date of publication of the international search report: 8 August 1996 (08.08.96)	
(54) Title: METHODS AND COMPOSITIONS RELATING TO PLANT Δ^6 PALMITOYL-ACYL CARRIER PROTEIN DESATURASE					
(57) Abstract <p>A plant Δ^6 palmitoyl-acyl carrier protein desaturase, the gene encoding the desaturase, and transgenic plants and plant cells containing the heterologous DNA encoding the desaturase are described. The desaturase introduces a double bond at the sixth carbon atom from the carboxyl end of a 16 carbon saturated fatty acid, and is therefore useful in production of plant seeds having a modified fatty acid profile.</p>					

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

INTERNATIONAL SEARCH REPORT

 Inter national Application No
 PC1/US 95/13784

 A. CLASSIFICATION OF SUBJECT MATTER
 IPC 6 C12N15/53 C12N15/82 C12N9/02 A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

 Minimum documentation searched (classification system followed by classification symbols)
 IPC 6 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 122, no. 13, 27 March 1995 Columbus, Ohio, US; abstract no. 156368, CAHOON, EDGAR BENJAMIN: "Synthesis and metabolism of .DELTA.6 monosaturated fatty acids in developing seed of Umbelliferae species and Thunbergia alata" XP002000650 see abstract & DISSERTATION (1994) 219 PP. AVAIL.: UNIV. MICROFILMS INT., ORDER NO. DA9431218 FROM: DISS. ABSTR. INT. B 1995, 55(7), 2482 . --- -/--	1,6-9,13

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"A" document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

16 April 1996

09.05.96

 Name and mailing address of the ISA
 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
 Fax (+ 31-70) 340-3016

Authorized officer

Maddox, A

INTERNATIONAL SEARCH REPORT

Inter- national Application No
PC1/US 95/13784

C. (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	J. BIOL. CHEM. (1994), 269(44), 27519-26, 4 November 1994, XP002000645 CAHOON, EDGAR B., ET AL.: ".DELTA.6 hexadecenoic acid is synthesized by the activity of a soluble .DELTA.6 palmitoyl-acyl carrier protein desaturase in Thunbergia alata endosperm" see the whole document ---	1,6-9,13
A	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 89, 1992, WASHINGTON US, pages 11184-11188, XP002000646 CAHOON, E.B., ET AL.: "Expression of a coriander desaturase results in petroselinic acid production in transgenic tobacco" see the whole document ---	1-19
A	PLANT PHYSIOLOGY, vol. 106, October 1994, pages 807-808, XP002000647 CAHOON, E.B., ET AL.: "cDNAs for isoforms of the delta9-stearyl-acyl carrier protein desaturase from Thunbergia alata endosperm" see the whole document ---	1-19
A	PLANT MOLECULAR BIOLOGY, vol. 26, October 1994, pages 631-642, XP002000648 SCHMIDT, H., ET AL.: "Purification and PCR-based cloning of a plastidial n-6 desaturase" see the whole document ---	1-19
A	LIPIDS, vol. 6, 1971, pages 712-714, XP002000649 SPENCER, G.F., ET AL.: "Occurrence of cis-6-hexadecenoic acid as the major component of Thunbergia alata seed oil" see the whole document ---	1-19
A	WO,A,93 11245 (DU PONT) 10 June 1993 see page 85 - page 88 ---	1-19
A	WO,A,94 01565 (MINI AGRICULTURE & FISHERIES) 20 January 1994 see the whole document -----	1-19

Form PCT/ISA/215 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International Application No.

PCI/US 95/13784

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
WO-A-9311245	10-06-93	AU-B-	3228893	28-06-93
		CA-A-	2124673	10-06-93
		EP-A-	0616644	28-09-94
		JP-T-	7501701	23-02-95

WO-A-9401565	20-01-94	AU-B-	4508393	31-01-94
		EP-A-	0603371	29-06-94
